

The Roaming Atom: Straying from the Reaction Path in Formaldehyde Decomposition

D. Townsend^{1,2}, S. A. Lahankar³, S. K. Lee^{1,2,3}, S. D. Chambreau³, A. G. Suits^{1,2,3*},
X. Zhang⁴, J. Rheinecker⁴, L. B. Harding⁵ and J. M. Bowman^{4*}

¹Department of Chemistry, Stony Brook University, Stony Brook, NY 11794

²Chemistry Department, Brookhaven National Laboratory, Upton NY 11973

³Department of Chemistry, Wayne State University, Detroit MI 48202

⁴Department of Chemistry and Cherry L. Emerson Center for Scientific Computation, Emory University, Atlanta, GA, 30322

⁵Chemistry Division, Argonne National Laboratory, Argonne IL 60439

[*asuits@wayne.edu](mailto:asuits@wayne.edu); jimbowma@emory.edu

We present a combined experimental and theoretical investigation of formaldehyde (H₂CO) dissociation to H₂ and CO at energies just above the threshold for competing H elimination. High-resolution state-resolved imaging measurements of the CO velocity distributions reveal two dissociation pathways. The first proceeds through a well-established transition state to produce rotationally excited CO and vibrationally cold H₂. The second dissociation pathway yields rotationally cold CO in conjunction with highly vibrationally excited H₂. Quasi-classical trajectory calculations performed on a global potential energy surface for H₂CO suggest that this second channel represents an intramolecular hydrogen abstraction mechanism: one hydrogen atom explores large regions of the potential energy surface before bonding with the second H atom, by-passing the saddle point entirely.